Draft version January 15, 2013
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1301.2814v1 [astro-ph.SR] 13 Jan 2013

# PSR J1840−1419: A VERY COOL NEUTRON STAR

E. F. Keane[1], M. A. McLaughlin[2], M. Kramer[1,3], B. W. Stappers[3], C. G. Bassa[3], M. B. Purver[3] & P. Weltevrede[3]

[1] Max Planck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany.
[2] Department of Physics, West Virginia University, Morgantown, WV 26506, USA.
[3] University of Manchester, Jodrell Bank Centre for Astrophysics, School of Physics & Astronomy, Manchester M13 9PL, UK.

*Draft version January 15, 2013*

## ABSTRACT

We present upper limits on the X-ray emission for three neutron stars. For PSR J1840−1419, with a characteristic age of 16.5 Myr, we calculate a blackbody temperature upper limit (at 99% confidence) of $kT_{\rm bb}^{\infty} < 24^{+17}_{-10}$ eV, making this one of the coolest neutron stars known. PSRs J1814−1744 and J1847−0130 are both high magnetic field pulsars, with inferred surface dipole magnetic field strengths of $5.5 \times 10^{13}$ and $9.4 \times 10^{13}$ G, respectively. Our temperature upper limits for these stars are $kT_{\rm bb}^{\infty} < 123^{+20}_{-33}$ eV and $kT_{\rm bb}^{\infty} < 115^{+16}_{-33}$ eV, showing that these high magnetic field pulsars are not significantly hotter than those with lower magnetic fields. Finally, we put these limits into context by summarizing all temperature measurements and limits for rotation-driven neutron stars.

*Subject headings:* pulsar: general — stars: magnetic fields — X-rays: stars

## 1. INTRODUCTION

Measurements of the temperatures of neutron stars (NSs) are few and far between. They are often impossible as, for ages $\gtrsim 1$ Myr, the star will have experienced significant cooling since its formation (Yakovlev & Pethick 2004). The thermal emission from NSs peaks in soft X-rays, below $\sim 1$ keV, and the luminosities are such that sources even a few kiloparsecs away are difficult to detect. Currently, a total of 37 measurements, and 9 upper-limit measurements, of rotation-driven NS temperatures have been published (Li et al. 2005; Caraveo et al. 2010; Abdo 2010; Speagle et al. 2011; Chang et al. 2011; Zhu et al. 2011; Marelli et al. 2011; Guver et al. 2012; Pires et al. 2012). Of these, only 11 are for NSs with characteristic ages older than 1 Myr (excluding 5 millisecond pulsars for whom we have no meaningful estimate of age), and only three of those greater than 10 Myr: PSRs B0950+08, J2144−3933 and J0108−1431. The emission from PSR B0950+08, a 17.5 Myr pulsar, is non-thermal, with a power-law spectrum, and an upper limit of $kT_{\rm bb}^{\infty} < 41$ eV, or $T < 0.48$ MK, on the thermal contribution (Becker et al. 2004). To put this in context, it is at the level of the lowest measured temperature for any NS: $kT_{\rm bb}^{\infty} = 43$ eV for Geminga (De Luca et al. 2005). PSR J2144−3933 is the slowest spinning pulsar known ($P = 8.5$ s) and an order of magnitude older (272 Myr) than PSR B0950+08 so that it is perhaps not surprising that no X-ray emission has been seen from it, with an upper limit of $kT_{\rm bb}^{\infty} < 20$ eV (Tiengo et al. 2011). Posselt et al. (2012) have recently reported the detection of PSR J0108−1431 (170 Myr) with $kT_{\rm bb}^{\infty} = 110$ eV, a source which seems to be undergoing a heating process.

In a recent re-analysis of the Parkes Multi-beam Pulsar Survey, PSR J1840−1419 was discovered (Keane et al. 2010). This 6.6-s pulsar shows sporadic radio emission with strong single pulses detected, on average, every $\sim 10$ rotation periods, at flux densities up to 1.7 Jy. At 16.5 Myr, PSR J1840−1419 has a very similar characteristic age to PSR B0950+08, and therefore might be expected to have somewhat similar X-ray properties. This, plus its relative proximity at 850 pc, make PSR J1840−1419 an excellent target for adding to our knowledge of the X-ray emission and temperature of old NSs. In this paper we present the result of a *Chandra* observation of PSR J1840−1419.

One factor which has been suggested to be a major factor in NS cooling is the magnetic field strength. Several authors have suggested that NSs with higher magnetic field strengths are hotter than those with lower field strengths, for the same age (Gonzalez et al. 2007; Zhu et al. 2009; Olausen et al. 2010; Zhu et al. 2011). Such an effect is theoretically predicted (Yakovlev & Pethick 2004) and the work of Geppert et al. (2004) and Pérez-Azorín et al. (2006) shows that high magnetic fields supress heat conductivity perpendicular to the field lines, naturally producing an anisotropic temperature distribution on the stellar surface with small hot regions at the magnetic poles. We examine this claim with *XMM-Newton* observations of two pulsars: J1814−1744 and J1847−0130, both of which have high magnetic field strengths, of $5.5 \times 10^{13}$ and $9.4 \times 10^{13}$ G, respectively.

Finally, we present an updated plot of $kT_{\rm bb}^{\infty}$ vs characteristic age for all X-ray measurements of, and upper limits on, rotation-powered NS temperature made thus far, and re-examine the temperature-magnetic field strength relationship.

## 2. OBSERVATIONS & RESULTS

### 2.1. *Calculating Limits*

For a non-detection, one could determine a count rate limit by following the procedure of Pivovaroff et al. (2000): (i) define an aperture on the image about the position of the source, and add up the counts in this region, $N$; (ii) define a background region of the same area (or appropriately scale to the same area) elsewhere on the image and add up the counts in the background region, $B$. The signal from the source is $N - B$ and the noise is $\sqrt{N+B}$. The signal-to-noise ratio is then

$(N - B)/\sqrt{N + B}$. To set a '$3 - \sigma$ limit' count rate we can then solve for $N$ in $9(N + B) = (N - B)^2$, where $B$ is known. We can see that in the $N \gg B$ 'photon-rich' case this tends to the expected Poisson value in the absence of background sky noise. However, in the 'photon-poor' case we clearly can not use this expression, e.g. for $B = 0.35$, $N = 10$ naïvely suggests a 99.7% detection/limit, however we know that the Poisson probability for 10 counts due to noise is $\approx 10^{-13}$%; for such a background rate 3 counts represents a 99.8% limit. The observations presented here are in the photon-poor scenario, so that using the Pivovaroff et al. (2000) procedure would result in a much poorer limit than what has truly been obtained.

To convert count rate limits to flux limits we need either some kind of absolute scale, e.g. from observing a standard source of known flux, or a model for our source, e.g. an assumed spectral form (McLaughlin et al. 2003). Below we calculate one set of limits for a blackbody model (whence we obtain temperature limits as a function of the emitting radius), and another set of limits for a non-thermal model with a power law with negative index of 2.0.

### 2.2. *PSR J1840−1419*

PSR J1840−1419 is an old pulsar (16.5 Myr) with a 6.6-s spin period. Due to its sporadic radio emission it was detected in a search for single pulses, rather than in a periodicity search (Keane et al. 2010). Pulsars discovered in this way are often referred to as "RRATs" (Keane & McLaughlin 2011). The properties of PSR J1840−1419, both measured and derived, are given in Table 1.

On 2011-02-20, we performed a 10-ks observation of PSR J1840−1419, using the ACIS-S detector on *Chandra* in the energy range $0.2 - 10.0$ keV. We detected only one photon within the $\sim 1$ arcsec (3 pixel) error circle of the position derived from pulsar timing (Keane et al. 2011). Assuming that this 1 count is consistent with background noise (the background rate being 0.24 counts/pixel), and ignoring the Poisson nature of the source, the count rate limit is $3 \times 10^{-4}$ counts/s (at 99% confidence).

In the case of a blackbody model this count rate limit implies a temperature limit of $kT^{\infty}_{\rm bb} < 24^{+17}_{-10}{\rm eV}(R^{\infty}_{\rm bb}/10\ {\rm km})^{-0.39}$, where the dependence on $R^{\infty}_{\rm bb}$ is a purely empirical fit to a simple power law of the form $T_{10}R^{\alpha}_{10}$, where $R_{10}$ is the blackbody radius in units of 10 km and $T_{10}$ is the temperature when $R_{10} = 1$: see top left panel of Figure 1. The error bars reflect both NE2001 'worst case' errors of a factor of two in the distance (Cordes & Lazio 2002) as well as factor of two uncertainties in the neutral hydrgoen column density (see below). The bottom right panel of Figure 1 shows all measured neutron star temperatures as a function of age. The PSR J1840−1419 limit is amongst the coolest of all published limits. The corresponding flux and luminosity limits are $f_{\rm bb} < 5.0 \times 10^{-14}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.61}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm bb} < 4.3 \times 10^{30}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.61}$ erg s$^{-1}$, respectively. In the case of the non-thermal model (power law, with negative index of 2.0), the flux and luminosity limits are $f_{\rm nt} < 3.0 \times 10^{-15}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm nt} < 2.6 \times 10^{29}$ erg s$^{-1}$, respectively.

The above estimates use a neutral hydrogen column density of $N_{\rm H} = 6 \times 10^{20}$ cm$^{-2}$ which is derived from the dispersion measure, assuming 10 neutral hydrogen atoms per free electron (Seon & Edelstein 1998). Predicted count levels were calculated using the standard PIMMS (Portable, Interactive Multi-Mission Simulator) tool[1].

### 2.3. *PSRs J1814−1744 & J1847−0130*

We also present the results of two *XMM-Newton* observations using the PN detector and medium filter in the energy range $0.15 - 15.0$ keV. PSR J1814−1744 was observed for 6.1 ks on 2004-10-21, and PSR J1847−0130 was observed for 17.0 ks on 2004-09-14. The observations of these young, high-B pulsars also resulted in non-detections. As above for PSR J1840−1419 we derive temperature upper limits for these sources of: $kT^{\infty}_{\rm bb} < 123^{+20}_{-34}{\rm eV}(R^{\infty}_{\rm bb}/10\ {\rm km})^{-0.23}$, for PSR J1814−0130, and $kT^{\infty}_{\rm bb} < 115^{+16}_{-33}{\rm eV}(R^{\infty}_{\rm bb}/10\ {\rm km})^{-0.21}$, for PSR J1847−0130 (see Figure 1). These limits are much less constraining given the much larger estimated distances (7.7 and 9.8 kpc respectively). The associated flux and luminosity limits for the blackbody scenario are: $f_{\rm bb} < 2.6 \times 10^{-13}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.77}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm bb} < 3.0 \times 10^{33}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.77}$ erg s$^{-1}$ for PSR J1814−1744; and $f_{\rm bb} < 3.2 \times 10^{-13}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.79}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm bb} < 2.3 \times 10^{33}(R^{\infty}_{\rm bb}/10\ {\rm km})^{1.79}$ erg s$^{-1}$ for PSR J1847−0130.

The non-thermal limits are: $f_{\rm nt} < 8.9 \times 10^{-15}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm nt} < 1.0 \times 10^{32}$ erg s$^{-1}$ for PSR J1814−1744; and: $f_{\rm nt} < 5.1 \times 10^{-15}$ erg s$^{-1}$ cm$^{-2}$ and $L^{\infty}_{\rm nt} < 3.6 \times 10^{31}$ erg s$^{-1}$ for PSR J1847−0130. For PSR J1814−1744 the value of $N_{\rm H}$ derived from the dispersion measure was $\sim 50$% higher than the maximum Galactic value for this line of sight, as calculated by the standard COLDEN tool[2], so the COLDEN value of $N_{\rm H} = 1.6 \times 10^{22}$ cm$^{-2}$ was used. For this reason $N_{\rm H}$ is unlikely to be underestimated, but we have allowed for the possibility that it may be overestimated by as much as a factor of two. Likewise, for PSR J1847−0130, the neutral hydrogen column density was derived to be $N_{\rm H} = 2.1 \times 10^{22}$ cm$^{-2}$. This value of $N_{\rm H}$ is very close to the maximum value for this line of sight, so that it too is unlikely to be an overestimate.

## 3. DISCUSSION & CONCLUSIONS

Although this paper reports three null results we deem it to be of the utmost importance to report such investigations to avoid duplicated efforts, wasted telescope resources, etc. Table 1 summarises the temperature, flux and luminosity limits for the sources reported here. From the non-thermal luminosity limits we can determine upper-limits on the non-thermal X-ray efficiency, $\eta = L_{\rm nt}/\dot{E}$, where $\dot{E}$ is the spin-down energy loss rate. Although the luminosity limits for the *XMM-Newton* observations are $\sim 2$ orders of magnitude poorer than for the *Chandra* observation of J1840−1419, the $\dot{E}$ values are correspondingly higher, such that for all 3 pulsars the limit turns out to be $\eta \lesssim 0.2$. Possenti et al. (2002) determined, for their study of 39 pulsars, that all sources had a value of $\eta$ less than $\eta_{\max} = 10^{-18.5}(\dot{E}/\mathrm{erg\ s^{-1}})^{0.48}$. Our $\eta$ limits are all significantly higher than this predicted value. However, we note this critical $\eta$ value is arrived at upon considering pulsars studied in the $2-10$ keV energy range, narrower than the observations reported here.

---

[1] http://asc.harvard.edu/toolkit/pimms.jsp

[2] http://asc.harvard.edu/toolkit/colden.jsp

These observations bring the total number of rotation-powered NSs with detections of thermal emission, or upper limits thereupon, to 49. In Figure 1 we show our limits, along with all previous measurements and limits. Zhu et al. (2011) suggest that there is a "hint" that the high magnetic field pulsars are hotter than the low magnetic field pulsars. To quantify this we first ignored the upper limits and divided the remaining data into 'low' and 'high' magnetic-field-strength sources, below and above an arbitrarily chosen field strength value of $10^{13}$ G. We also excluded three 'low' magnetic-field-strength sources as their fitted blackbody radii are so much smaller than those of the rest of the sources (at 33, 43 and 120 m) so that their thermal emission is thought to be due to some heating process (Misanovic et al. 2008). Comparing the two distributions with a Kolmogorov-Smirnov (K-S) test we find a K-S statistic of $D = 0.33$, and thus a probability of 0.31 that these two distributions are the same. There is therefore no evidence that the high magnetic field pulsars are hotter than those with lower magnetic fields. However we note the caveat that, in addition to the age estimates being uncertain (as mentioned above), the magnetic field estimates are uncertain, and should only be considered accurate to within an order of magnitude. Knowledge of the inclination angles between the magnetic and rotation axes would be needed to estimate the magnetic field strengths more accurately (e.g. as in Spitkovsky (2006)).

## ACKNOWLEDGMENTS

This work has been supported by NASA CXO guest observer support grant GO1-12059X, and made use of software provided by the Chandra X-ray Center. EFK thanks the FSM for support, K. J. Lee for useful discussions on statistics, R. P. Eatough for useful comments on the manuscript and the anonymous referee for providing valuable input that improved the quality of this paper. MAM is supported by the Research Corporation for Scientific Advancement.

## REFERENCES

Abdo A. A. e. a., 2010, ApJ, 712, 1209

Becker W., Weisskopf M. C., Tennant A. F., Jessner A., Dyks J., Harding A. K., Zhang S. N., 2004, ApJ, 615, 908

Caraveo P. A., De Luca A., Marelli M., Bignami G. F., Ray P. S., Saz Parkinson P. M., Kanbach G., 2010, ApJ, 725, L6

Chang C., Pavlov G. G., Kargaltsev O., Shibanov Y. A., 2011, ApJ, in press

Cordes J. M., Lazio T. J. W., preprint (arXiv:astro-ph/0207156)

De Luca A., Caraveo P. A., Mereghetti S., Negroni M., Bignami G. F., 2005, ApJ, 623, 1051

Geppert U., Küker M., Page D., 2004, A&A, 426, 267

Gonzalez M. E., Kaspi V. M., Camilo F., Gaensler B. M., Pivovaroff M. J., 2007, Astrophys. Space Sci., 36

Guver T., Gogus E., Ozel F., 2012, MNRAS, in press

Keane E. F., Kramer M., Lyne A. G., Stappers B. W., McLaughlin M. A., 2011, MNRAS, 838

Keane E. F., Ludovici D. A., Eatough R. P., Kramer M., Lyne A. G., McLaughlin M. A., Stappers B. W., 2010, MNRAS, 401, 1057

Keane E. F., McLaughlin M. A., 2011, Bulletin of the Astronomical Society of India, 39, 333

Li X. H., Lu F. J., Li T. P., 2005, ApJ, 628, 931

Marelli M., De Luca A., Caraveo P. A., 2011, ApJ, 733, 82

McLaughlin M. A. et al., 2003, ApJ, 591, L135

Misanovic Z., Pavlov G. G., Garmire G. P., 2008, ApJ, 685, 1129

Olausen S. A., Kaspi V. M., Lyne A. G., Kramer M., 2010, ApJ, 725, 985

Pérez-Azorín J. F., Miralles J. A., Pons J. A., 2006, A&A, 451, 1009

Pires A. M., Motch C., Turolla R., Schwope A., Pilia M., Treves A., Popov S. B., Janot-Pacheco E., 2012, A&A, submitted

Pivovaroff M., Kaspi V. M., Camilo F., 2000, ApJ, 535, 379

Posselt B., Arumugasamy P., Pavlov G. G., Manchester R. N., Shannon R. M., Kargaltsev O., 2012, ApJ, in press

Possenti A., Cerutti R., Colpi M., Mereghetti S., 2002, A&A, 387, 993

Seon K.-I., Edelstein J., 1998, AJ, 115, 2097

Speagle J. S., Kaplan D. L., van Kerkwijk M. H., 2011, ApJ, 743, 183

Spitkovsky A., 2006, ApJ, 648, L51

Tiengo A., Mignani R. P., de Luca A., Esposito P., Pellizzoni A., Mereghetti S., 2011, MNRAS, 412, L73

Yakovlev D. G., Pethick C. J., 2004, Ann. Rev. Astr. Ap., 42, 169

Zhu W., Kaspi V. M., Gonzalez M. E., Lyne A. G., 2009, ApJ, 704, 1321

Zhu W. W., Kaspi V. M., McLaughlin M. A., Pavlov G. G., Ng C.-Y., Manchester R. N., Gaensler B. M., Woods P. M., 2011, ApJ, 734, 44

**Table 1**
Measured and derived properties, and limits for 3 pulsars.

| Quantity | J1840−1419 | J1814−1744 | J1847−0130 |
|---|---|---|---|
| $P$ (s) | 6.6 | 4.0 | 6.7 |
| $\dot{P}$ (fs/s) | 6.3 | 745 | 1270 |
| $\tau_{\rm c}$ (kyr) | 16500 | 85 | 83 |
| $B$ ($10^{12}$ G) | 6.5 | 55.1 | 93.6 |
| $\dot{E}$ ($10^{30}$ erg s$^{-1}$) | 1.0 | 468 | 167 |
| $DM$ (cm$^{-3}$ pc) | 19 | 792 | 667 |
| $D$ (kpc) | 0.9 | 9.8 | 7.7 |
| Telescope | Chandra | XMM | XMM |
| Instrument | ACIS-S | PN | PN |
| Date | 2011-02-20 | 2004-10-21 | 2004-09-14 |
| Energy Range (keV) | $0.2 - 10$ | $0.15 - 15$ | $0.15 - 15$ |
| $T_{\rm obs}$ (ks) | 10.0 | 6.1 | 17.0 |
| $N_{\rm H}$ ($10^{20}$ cm$^{-2}$) | 6 | 157 | 205 |
| $kT^{\infty}_{\rm bb}$ (eV)† | $< 24^{+17}_{-10}$ | $< 123^{+20}_{-34}$ | $< 115^{+16}_{-33}$ |
| $T^{\infty}_{\rm bb}$ (MK)† | $< 0.28^{+0.19}_{-0.12}$ | $< 1.42^{+0.22}_{-0.39}$ | $< 1.33^{+0.19}_{-0.38}$ |
| $f_{\rm bb}$ ($10^{-14}$ erg cm$^{-2}$ s$^{-1}$)‡ | $< 5.0$ | $< 6.1$ | $< 8.9$ |
| $L_{\rm bb}$ ($10^{30}$ erg s$^{-1}$)‡ | $< 4.3$ | $< 692$ | $< 640$ |
| $f_{\rm nt}$ ($10^{15}$ erg cm$^{-2}$ s$^{-1}$) | $< 3.0$ | $< 8.9$ | $< 5.1$ |
| $L_{\rm nt}$ ($10^{30}$ erg s$^{-1}$) | $< 0.26$ | $< 102$ | $< 36$ |
| $\eta = L_{\rm nt}/\dot{E}$ | $< 0.26$ | $< 0.22$ | $< 0.22$ |

**Note**. — The measured and derived properties for the three pulsars discussed in this paper. The spin period, $P$, and its derivative, $\dot{P}$, are derived from pulsar timing techniques, and $\tau$, $B$ and $\dot{E}$, the characteristic age, magnetic field strength and spin-down energy respectively, are inferred from these (Keane et al. 2011). The distance, $D$, is derived from the measured $DM$ using the NE2001 model for the Galactic free electron distribution (Cordes & Lazio 2002). The 99%-confidence temperature limits are derived as outlined in the text. Non-thermal flux limits are given for a power law with negative index of 2.0. † these limits are for an emitting radius of 10 km, and scale as $(R^{\infty}_{bb}/10\ {\rm km})^{\alpha}$, where $\alpha$ equals $-0.39$, $-0.23$ and $-0.21$, respectively for the three sources (see main text). ‡ The blackbody flux and luminosity limits similarly scale as $(R^{\infty}_{bb}/10\ {\rm km})^{2-\alpha}$.



**Figure 1.** The first three plots show the temperature limit (i.e. which produces the observed count rate limit for an assumed blackbody model) as a function of the assumed emitting radius, for the *Chandra* observation of PSR J1840−1419 (10 ks) and the *XMM-Newton* observations of PSRs J1814−1744 (6 ks) and J1847−0130 (17 ks). The points are fit with a simple power-law of the form $T_{10}(R_{10})^{\alpha}$. In each case the middle curve is for the nominal dispersion-measure-derived distance. The cooler and hotter curves are for 'worst-case' distance errors of a factor of 2 (Cordes & Lazio 2002). For clarity of presentation we show here only the curves for the nominal $N_{\rm H}$ values quoted in Table 1. The fourth plot shows all known temperature values as a function of characteristic age. Sources with inferred magnetic field strengths $\geq 10^{13}$ G are marked in blue, whereas those with lower values are marked in black. Open circles denote upper limits. The three upper limit presented in this paper are marked with large circles for clarity. Note that the symbols for PSRs J1814−1744 and J1847−0130 are largely overlapping as their ages and derived limits are very similar.